



Cambridge IGCSE™

CANDIDATE
NAME

| |
|--|
| |
|--|

CENTRE
NUMBER

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

CANDIDATE
NUMBER

| | | | |
|--|--|--|--|
| | | | |
|--|--|--|--|



ADDITIONAL MATHEMATICS

0606/02

Paper 2

For examination from 2025

SPECIMEN PAPER

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a scientific calculator where appropriate.
- You must show all necessary working clearly.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- For π , use either your calculator value or 3.142.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

List of formulas

Equation of a circle with centre (a, b) and radius r .

$$(x - a)^2 + (y - b)^2 = r^2$$

Curved surface area, A , of cone of radius r , sloping edge l .

$$A = \pi r l$$

Surface area, A , of sphere of radius r .

$$A = 4\pi r^2$$

Volume, V , of pyramid or cone, base area A , height h .

$$V = \frac{1}{3}Ah$$

Volume, V , of sphere of radius r .

$$V = \frac{4}{3}\pi r^3$$

Quadratic equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial theorem

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series

$$u_n = a + (n - 1)d$$

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n - 1)d\}$$

Geometric series

$$u_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1 - r} \quad (|r| < 1)$$

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulas for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$


$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}ab \sin C$$

1 (a) Solve the equation $5|5x - 7| - 1 = 14$.

[3]

(b)



The diagram shows the graph of $y = f(x)$, where $f(x) = 2(x + 1)^2(x - 1)$.

Use the graph to solve the inequality $f(x) \leq -1$.

[3]

- 2 For variables x and y , plotting $\ln y$ against $\ln x$ gives a straight-line graph passing through the points $(6, 5)$ and $(8, 9)$.

Show that $y = e^p x^q$ where p and q are integers to be found. [4]

- 3 Find the values of the constant k for which the equation $(2k - 1)x^2 + 6x + k + 1 = 0$ has real roots. [5]

4 A photographer takes 12 different photographs. There are 3 photographs of sunsets, 4 of oceans and 5 of mountains.

(a) The photographs are arranged in a line on a wall.

(i) Find the number of possible arrangements if the first photograph is of a sunset and the last photograph is of an ocean. [2]

(ii) Find the number of possible arrangements if all the photographs of mountains are next to each other. [2]

(b) Three of the photographs are selected for a competition.

(i) Find the number of different possible selections if no photograph of a sunset is chosen. [2]

(ii) Find the number of different possible selections if one photograph of each type (sunset, ocean, mountain) is chosen. [2]

- 5 Given that $y = \tan x$, use calculus to find the approximate change in y as x increases from $-\frac{\pi}{4}$ to $h - \frac{\pi}{4}$, where h is small. [3]

- 6 A curve has equation $y = \ln(5 - 3x)$ where $x < \frac{5}{3}$. The normal to the curve at the point where $x = -5$, cuts the x -axis, at the point P .

Find the equation of the normal and the x -coordinate of P . [7]

7 Solutions to this question by accurate drawing will not be accepted.

A circle has equation $x^2 + y^2 - 16x - 10y + 73 = 0$.

(a) (i) Find the coordinates of the centre of the circle and the length of the radius. [3]

(ii) Hence show that the point $(10, 6.5)$ lies inside the circle. [2]


(b) A different circle has equation $(x - 10)^2 + (y - 6.5)^2 = 2.25$.

Show that the two circles touch. You are not required to find the coordinates of the common point. [1]

8 (a) (i) Show that $\frac{\cos^2 2x}{1 + \sin 2x} = 1 - \sin 2x$. [2]

(ii) Hence solve the equation for $\frac{3 \cos^2 2x}{1 + \sin 2x} = 1$ for $0^\circ \leq x \leq 90^\circ$. [4]

(b) Solve the equation $\cot\left(y - \frac{\pi}{2}\right) = \sqrt{3}$, where y is in radians and $0 \leq y \leq \pi$. [3]



The diagram shows a rectangle $ABCD$ and an arc AXB of a circle with centre at O , the midpoint of DC . The length of BC is 8 cm and the length of DC is 30 cm. Angle AOB is θ radians.

(a) Find the perimeter of the shape $ADOCBX$. [5]

(b) Find the area of the shape $ADOCBX$. [2]

- 10 (a) In the expansion of $\left(2k - \frac{x}{k}\right)^5$, where k is a constant, the coefficient of x^2 is 160.

Find the value of k .

[3]


- (b) (i) Find the first 3 terms in the expansion of $(1 + 3x)^6$, in ascending powers of x . Simplify the coefficient of each term.

[2]

- (ii) When the expansion of $(1 + 3x)^6(a + x)^2$ is written in ascending powers of x , the first three terms are $4 + 68x + bx^2$, where a and b are constants.

Find the value of a and the value of b . [3]

11 In this question, all lengths are in centimetres.




The diagram shows a cone of base radius x , height y and sloping edge $\sqrt{x^2 + y^2}$. The volume of the cone is $10\pi \text{ cm}^3$.

(a) Show that the curved surface area, S , of the cone is given by $S = \frac{\pi\sqrt{x^6 + 900}}{x}$. [3]

- (b) Given that x can vary and that S has a minimum value, find the value of x for which S is a minimum. [5]

12



The diagram shows part of the curve $y = (9-x)(x-3)$ and the line $y = k-3$, where $k > 3$. The line through the maximum point of the curve, parallel to the y -axis, meets the x -axis at A . The curve meets the x -axis at B , and the line $y = k-3$ meets the curve at the point $C(k, k-3)$.

Find the area of the shaded region.

[9]

Continuation of working space for Question 12.

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge University Press & Assessment. Cambridge University Press & Assessment is a department of the University of Cambridge.