



Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

PHYSICAL SCIENCE

0652/31

Paper 3 Theory (Core)

October/November 2024

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has 24 pages. Any blank pages are indicated.



1 (a) A list of atoms and ions is shown.

Al C Ca²⁺ Cl⁻ Cu Fe K⁺ Mg

2

Use the symbols of the atoms or ions to answer the questions that follow.

Each symbol may be used once, more than once or not at all.

State which atom or ion:

(i)	is used in food containers because of its resistance to corrosion	
		[1]
(ii)	is extracted from the ore bauxite by electrolysis	
		[1]
(iii)	burns with a lilac flame	
		[1]
(iv)	is a non-metal with two electron shells	-4-
, ,		[1]
(v)	has the same number of electrons in its outer shell as argon	[41
(s.:i)	is a motal that is loss reactive than hydrogen	[1]
(vi)	is a metal that is less reactive than hydrogen.	[4]
		ניו

(b) Ammonia, NH_3 , is a covalent molecule.

Complete Fig. 1.1 to show the dot-and-cross diagram for a molecule of NH_3 .

Show outer shell electrons only.




Fig. 1.1

© UCLES 2024

 * 00800000003	

1	c)	Describe a	test for	ammonia d	as and	state the	observation	if	ammonia i	is	nresent
V.	\sim	Dodding a	toot ioi	arriirioriia g	ao ana	otato tilo		•••	arriirioriia i	0	produit.

test	 	
observation	 	
	 	 oı

[Total: 10]

[2]



2 A cyclist starts from rest.

She travels with constant acceleration for $5.0\,\mathrm{s}$ and reaches a speed of $10\,\mathrm{m/s}$. She travels a distance of $25\,\mathrm{m}$ while accelerating.

The cyclist continues at 10 m/s for a further 30 s.

(a) Calculate the distance the cyclist travels while moving at 10 m/s for 30 s.

(b) On Fig. 2.1, plot a speed–time graph of the motion of the cyclist.




Fig. 2.1

(c) On your graph in Fig. 2.1, shade the area which represents the 25 m the cyclist travels while accelerating. [1]



(d) Another cyclist travels 50 m in 10 s while accelerating.

He travels a further 400 m in 50 s at constant speed.

5

Calculate the average speed of the cyclist.

[Total: 7]



BLANK PAGE



(b)



(a) Fig. 3.1 shows an isotope of silicon.

³⁰Si

7

Fig. 3.1

	Table 3 1	
Cor	mplete Table 3.1.	
		[1]
(iii)	Determine the number of neutrons in the isotope in Fig. 3.1.	
		[1]
(ii)	Determine the number of protons in the isotope in Fig. 3.1.	
		[2]
(i)	Define the term isotopes.	

particle	relative charge	relative mass
proton		
neutron		
electron		

[3]

[1]

- (c) Silicon is used with aluminium to form an alloy.
 - Circle the term that best describes an alloy.

element compound mixture solvent pure

(ii) Complete the sentence.

> An alloy of aluminium is used instead of pure aluminium because the properties of the [1] alloy are

> > [Total: 9]

A crane lifts a large bucket containing concrete. The volume of the concrete is 2.0 m³. The density of the concrete is 2600 kg/m³.

Calculate the mass of the concrete.

mass	=	ka	[2]
111433		Kg	L - .

(b) Calculate the weight of a 55 kg concrete block.

 $[g = 10 \,\text{N/kg}]$

(c) The energy supplied to the electric motor of the crane to lift a load is 20 000 J.

The gravitational potential energy of the load increases by 14000 J.

Explain how this information indicates that the crane is **not** 100% efficient.

(ii) Circle the phrase in the list which correctly completes the sentence.

more energy more force more power more resistance

To lift the same load in less time, the electric motor of the crane needs

[1]


DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN



(d) Fig. 4.1 shows a crane which moves a load (L) from P to Q.

The crane has a counterweight (C).



9

Fig. 4.1

The load is moved from P to Q.

Counterweight **C** is moved to the right to stop the crane falling over.

Explain in terms of moments why the counterweight is moved to the right.

explanation	 	
•		
		 [2]
	 	 [2]

[Total: 7]



5 Fig. 5.1 shows a microwave oven with food inside.

A microwave oven uses microwaves to cook food.




Fig. 5.1

a)	(i)	Name the type of electromagnetic radiation which passes through the window so y can see the food cooking.	′ou
			[1]
	(ii)	The food is too hot to eat and is allowed to cool for several minutes.	
		Name the type of electromagnetic radiation which is emitted by the food as it cools.	
			[1]
b)		me hot food is put onto a cold plate. veral minutes later the plate is hot.	
	Sta	te and describe the process which causes the plate to become hot.	
			[2]

(ii)



(c) (i) Fig. 5.2 shows a liquid-in-glass thermometer.




Fig. 5.2


Explain why the level of the liquid in the narrow tube changes when the thermometer is put in hot food.
[2]
Another thermometer has a smaller reservoir of liquid than shown in Fig. 5.2. The diameter of the narrow tube is the same as shown in Fig. 5.2. This thermometer is put into the same hot food.
Predict the difference, if any, in the level of liquid in the narrow tube of this thermometer.
[1]
[Total: 7]

BLANK PAGE





(a) Fig. 6.1 shows the apparatus used for the electrolysis of an aqueous solution.



13

Fig. 6.1

	•							
(i)	State the names of X , Y and Z .							
	X							
	Υ							
	Z			[0]				
				[2]				
(ii)	Fig. 6.1 shows the electrolysis of an	aqueous solutior	1.					
	Circle the other type of substance th	at can be electro	lysed.					
	gas mol	ten	solid					
				[1]				
(iii)	Name a suitable inert material to use	for X and Y .						
				[1]				
A g	A gas is produced during the electrolysis shown in Fig. 6.1.							
Des	escribe the observation that shows a ga	s is produced.						
				[1]				

[Total: 5]

(b)

Fig. 7.1 shows an electromagnet. It consists of a core inside a coil made from turns of wire.

14





Fig. 7.1

(i)	Circle one	material from	the list which	makes the best	core for the	electromagne
-----	------------	---------------	----------------	----------------	--------------	--------------

	air	aluminium	soft iron	steel	zinc	[1]
(ii)	The best core	e for the electromag	net is chosen.			
	State two ch	anges that increase	the strongth of the	magnatia fia	ld of the electrome	anot

[2]

(b) The electromagnet is used to make a loudspeaker, as shown in Fig. 7.2.



15

Fig. 7.2

A bar magnet is attached to a card.

The card hangs from a spring.

The electromagnet is a small distance below the bar magnet.

(i)	There is a small current in the electromagnet. The card moves down a short distance.
	Suggest how the card moves when the current in the electromagnet is:
	increased

(ii) A person hears the sound from the loudspeaker in Fig. 7.2.

Put the statements in the correct order, 1 to 4, to describe how the person hears the sound.

in the opposite direction.

	order
The sound wave makes the ear drum vibrate.	
The card makes the air vibrate.	
The electromagnet makes the bar magnet and card move up and down.	
The vibrations in the air travel as a sound wave.	

(c) State the lowest frequency that is audible to a healthy human ear.



[Total: 7]

[2]

[1]

[Turn over

8 (a) Fig. 8.1 shows an apparatus used in an industrial process.




Fig. 8.1

(i)	The raw material labelled A is a mixture of hydrocarbons.	
	Explain what is meant by hydrocarbon.	
		[2]
(ii)	Name the raw material labelled A .	
		[1]
(iii)	Tick (✓) the process that happens in B .	
	cooling	
	decomposition	
	dissolving	
	heating	
		[1]
(iv)	Name the method of separation that happens in C .	
		[1]
(v)	State one use for bitumen.	
		[1]

(c)




(b) Gasoline is a carbon-containing substance. Gasoline is a fuel.

Explain why the complete combustion of gasoline is a concern.
[2]
Name a fuel that does not contain carbon.
[1]
[Total: 9]

[2]

9 (a) Fig. 9.1 shows two plastic rods, A and B.



18

Fig. 9.1

Rod A is on a balance.

Rod B is moved closer to rod A.

Complete Table 9.1 to show how different charges on rods **A** and **B** change the reading on the balance when rod **B** is moved closer to rod **A**.

Only use the words decrease, increase, unchanged.

Table 9.1

charge on rod A	charge on rod B	change in reading on the balance
neutral	neutral	
positive	negative	
negative	positive	
positive	positive	

(b)	A plastic rod is ru	bed with a cloth.	The rod becomes	negatively	charged
-----	---------------------	-------------------	-----------------	------------	---------

Explain in terms of particles now the rod becomes negatively charged.
[2]
A metal rod is rubbed with a cloth.
Suggest why the metal rod does not become charged

[Total: 5]



(c)



10 Iron is a metal.

(a)	(i)	State the conditions needed for iron to rust.	
			[2]
	(ii)	State a barrier method which prevents rusting.	
			[1]
(b)	Fig.	10.1 shows the order of metals in the reactivity series.	
	Writ	te an X to show the position of iron in the reactivity series.	
		potassium	
		sodium	
		calcium	
		magnesium	
		aluminium	
		zinc	
		copper	
		Fig. 10.1	[1]
(c)	Writ	te a word equation for the reaction of iron with dilute hydrochloric acid.	
			[2]
(d)	Stee	el is made from iron.	
	Stat	te one use for mild steel.	
			[1]
		[Total	: 7]

(a)	Son	ne nuclei undergo radioactive decay.	
	Stat	e what is meant by radioactive decay.	
			[2]
(b)	An i	sotope of radon has a half-life of 2 hours.	
	Tick	(<) the two correct statements about the nuclei in a sample of this isotope of radon.	
	All t	he nuclei decay in 2 hours.	
	All t	he nuclei decay in 4 hours.	
	Any	one nucleus can decay at any time.	
	Half	the nuclei decay in 2 hours.	
			[2]
(c)	Alpł	na-particles, beta-particles and gamma-rays have different penetrating abilities.	
	(i)	Name a material that absorbs alpha-particles but not beta-particles.	
			[1]
	(ii)	Name a material that absorbs alpha-particles and beta-particles but not gamma-rays	
			[1]

[Total: 7]



(d) Name one method of detecting ionising radiation.



BLANK PAGE



BLANK PAGE



* 0000800000023 *

23

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.





The Periodic Table of Elements

				• •																
	III/	2 Helium	t (Ne S	neon 20	18	Ar	argon 40	36	궃	krypton 84	55	Xe	xenon 131	98	R	radon -	118	Og	oganesson
			σ	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Н	iodine 127	85	Ą	astatine -	117	<u>S</u>	tennessine -
			α	· O	oxygen 16	16	S	sulfur 32	34	Se	selenium 79	52	<u>L</u>	tellurium 128	84	Ро	molouium –	116	_	livermorium -
	>		7	Z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	Ξ	bismuth 209	115	Mc	moscovium -
	≥		œ	· O	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pp	lead 207	114	ŀΙ	flerovium -
	≡		ĸ	. ш	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	11	thallium 204	113	R	nihonium –
									30	Zn	zinc 65	48	р О	cadmium 112	80	Hg	mercury 201	112	C	copemicium
									59	Cn	copper 64	47	Ag	silver 108	6/	Au	gold 197	111	Rg	roentgenium -
Group									28	Z	nickel 59	46	Pq	palladium 106	78	₹	platinum 195	110	Ds	damstadtium -
ي			_						27	ပိ	cobalt 59	45	몬	rhodium 103	77	i	iridium 192	109	Ψ	meitnerium -
		T hydrogen	-						26	Ьe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	H	hassium
									25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium
			_	loq	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
		X	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	<u>Б</u>	tantalum 181	105	Dp	dubnium –
				atc	re				22	F	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	꿒	rutherfordium -
									21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=		4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	S	strontium 88	99	Ba	barium 137	88	Ra	radium
	_		ď	· '⊐	lithium 7	1	Na	sodium 23	19	×	potassium 39	37	Вb	rubidium 85	55	S	caesium 133	87	μ̈	francium

24

57 58 59 60 61 62 63 66 67 68 69 70 71 La Ce Pr Nd Pm Smartin Eu Gd Tb Dy HO Er Tm Yb Lu 139 140 141 144 - 150 152 157 159 163 167 169 173 175 89 90 91 92 94 95 96 97 98 99 100 101 102 133 Ac Th Pa Wp Pu Am Cm Bk Cf ES Fm Md No Lr 322 231 238 -							
58 59 60 61 62 63 66 66 67 68 69 Ce Pr Nd Pm Sm Eu Gd Tb Dy HO Er Tm certum prosecodymium recodymium semaritum europium gadolinium tenbium dysprosium holmium erbium thullium promium erbium thullium	71		lutetium 175	103	בֿ	lawrencium	I
58 59 60 61 62 63 64 65 66 67 68 Certum certum 140 Preseodymium 141 Promethium 	70	Υp	ytterbium 173	102	2	nobelium	I
Ce Pr Nd Pm Sm Eu Gd Tb Dy Ho certum pressedymium promethium samarium samarium Europium Gd Tb Dy Ho 140 141 144 - 150 157 157 159 163 165 90 91 92 93 94 95 96 97 98 99 Infortum uranium uranium plutonium plutonium americium curium berkelium crificinium enfortium	69	TB	thulium	101	Md	mendelevium	ı
Ce Pr Nd Pm Sm Eu Gd Tb Dy certum prosecdymium promethium samarium samarium europium gadolinium tentum dysprosium 140 141 144 - 150 157 159 163 90 91 92 93 94 95 96 97 98 Th Pa Np Pu Am Cm Bk Cf thorium uranium uranium plutonium americum curium berkelium carifornium	89	Щ	erbium 167	100	Fm	fermium	ı
Ce Pr Nd Pm Sm Eu Gd Tb Cerium praseodymium promethium samarium europium gadolinium terbium cerium 140 141 144 - 150 157 159 90 91 92 93 94 95 96 97 Th Pa Np Pu Am Cm BK thorium protectinium uranium uranium pertettium pertettium 232 231 238 - - - -	29	웃	holmium 165	60	Es	einsteinium	ı
58 59 60 61 62 63 64 Ce Pr Nd Pm Sm Eu Gd cerium praseodymium reodymium prometrium samarium europium gdodinium 140 141 144 - 150 152 157 90 91 92 93 94 95 96 Th Pa U Np Pu Am Cm thorium protectinium uranium reprunium putronium americum curium 232 231 238 - - - -	99	ρ	dysprosium	2 8	ŭ	californium	I
S8 59 60 61 62 63 Ce Pr Nd Pm Sm Eu certum praseodymlum neodymlum promethium samarium europium 140 141 144 - 150 152 90 91 92 93 94 95 Th Pa U Np Pu Am thorium protecthium uranium nepturium piutonium americium 232 231 238 - - -	99	Д	terbium 150	26	ă	berkelium	I
S8 59 60 61 62 Ce Pr Nd Pm Sm certum praseodymium neodymium promethium samartum 140 141 144 - 150 90 91 92 93 94 Th Pa U Np Pu thorium protectinium urranium nepturium plutonium 232 231 238 - - -	64	Вg	gadolinium 157	96	S	curium	I
Ce Pr Nd Pm seadymium promethium seadymium promethium seadymium neodymium seadymium seadymium promethium seadymium seadymium seadymium seadymium seadymium seadymium promethium promethium promethium seadymium seadymiu	63	Ē	europium 152	95	Am	americium	I
Ce Pr Nd certum praseodymium reodymium 140 141 144 90 91 92 Th Pa U thorium protectinum 232 231 238	62	Sm	samarium 150	94	Pu	plutonium	ı
Ce Pr cerium praseodymium n 140 91 Th Pa thorium protactinium 232 231	61	Pm	promethium	03	a N	neptunium	ı
Ce certum 140 90 90 Th thortum 232	09	PΝ	neodymium 144	6	: D	uranium	238
	29	P	praseodymium 1.4.1	6	Ра	protactinium	231
La lanthanum 139 89 AC actinium	58	Ce	cerium 140	06	H H	thorium	232
	22	Га	lanthanum 130	82	g S	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).